

ORIGINAL ARTICLE

OPEN ACCESS

Pak. J. Adv. Med. Med. Res.

Outcomes Of Functional Endoscopic Sinus Surgery (FESS) In Patients With Chronic Rhino sinusitis With Nasal Polyps

Tahreem Fatima¹, Shaker Ullah², Imran Khan³, Mushtaq Ahmad⁴, Muhammad Afaq⁵, Osama Nawaz⁶, Wasigh Ali⁷

¹Registrar ENT Department Khyber Teaching Hospital Peshawar

²Assistant Professor ENT Department Khyber Teaching Hospital Peshawar

^{1,4,5,6,7}Post graduate Trainees in ENT Department Khyber Teaching Hospital Peshawar

ABSTRACT

Background: Chronic rhino sinusitis with nasal polyps (CRS with NP) is a chronic inflammatory condition characterized by nasal congestion, olfactory dysfunction, and a diminished quality of life. Functional endoscopic sinus surgery (FESS) has emerged as a preferred treatment option for patients who do not respond adequately to medical management. While FESS is associated with improvements in symptom control and quality of life, the recurrence of polyps remains a significant concern, along with varying treatment outcomes. Therefore, it is crucial to assess the effectiveness and post-surgical outcomes of FESS in CRS with NP to optimize patient management strategies.

Objectives: To evaluate clinical outcomes, symptom relief, and quality-of-life improvement after FESS in patients with (CRS with NP) and discuss the results in relation to factors associated with recurrence and postoperative recovery.

Study design: A Cross-sectional study.

Place and Duration of study: Department of ENT Khyber Teaching Hospital Peshawar From January 2023 to January 2024

Methods: A Cross-sectional study was performed on patients diagnosed with (CRS with NP) FESS. Systems preoperative assessment was performed by symptom scoring with SNOT-22 and endoscopic probing. Follow-up was performed postoperatively after 3, 6 and 12 months with an evaluation of the symptomatic improvement, recurrences, and complications. SPSS was used to analyze the data and continuous variables were reported as mean + standard deviation and comparisons tested by paired t-tests. Statistical significance was predetermined to be $p < 0.05$.

Results: 75 patients with the mean age of 44.7 ± 12.5 years were involved in the study. Postoperative results showed a great improvement in SNOT-22 scores, which dropped to 28.7 (7.6) at 12 months ($p < 0.001$) compared to preoperative values of 62.4 (9.8). Nasal congestion resolved in 92 percent of the patients, hyposmia in 63 percent and headache in 62 percent. Recurrence was in 24 percent of the cases. Major complications were not reported. Better symptomatic outcome was observed with extended FESS based on symptomatic relief compared to limited FESS ($p = 0.028$). The quality-of-life scores showed better results which signify that FESS is effective in the reduction of symptoms and increases in day-to-day functions in (CRS with NP) patients.

Conclusion: Functional Endoscopic Sinus surgery markedly relieves the symptoms, quality of life and olfactory process in patients of chronic rhino sinusitis with nasal polyps. Recurrences are problematic; however, thorough patient selection and prolonged incision are options to improve the prognosis. FESS is safe with low complication rates, and offers sustained symptomatic relief, which supports its position as the surgery of first choice in a patient with medically refractory (CRS with NP).

Keywords: FESS, CRWNP, Outcomes, Recurrence

How To Cite: Fatima T, Ullah S, Khan I, Ahmad M, Afaq M, Nawaz O, Ali W. Outcomes of functional endoscopic sinus surgery (FESS) in patients with chronic rhino sinusitis with nasal polyps: original article. Pak J Adv Med Med Res. 2023;3(2):118-123; <https://doi.org/10.69837/pjammr.v3i2.78>

Corresponding Author: Imran Khan

Assistant Professor ENT Department Khyber Teaching Hospital
Peshawar

Email: imranam@hotmail.com

ORCID: <https://orcid.org/0000-0003-3152-2189>

Cell No: +92 344 8998856

OJS- Article Tracking

Received	January	24-2025
Revised	March	16-2025
Accepted	May	29 -2025
Published	July	10- 2025

INTRODUCTION

Chronic rhino sinusitis with nasal polyps (CRWNP) is a well-known condition of the Sino nasal mucosal inflammation with persistent symptoms of blockage of nostrils, loss of smell, pressure in the face and postnasal drip [1]. Pathogenesis is a complicated interaction of genetic predisposition, exposure to environments, immunological imbalance resulting in mucosal inflammation and formation of polyps [2]. Medical options such as intranasal corticosteroids, saline nose-washes, and systemic treatments find little relief especially in severe cases and in those with other diseases such as asthma [3]. Functional Endoscopic Sinus Surgery (FESS) has become a pillar in the treatment of (CRWNP) to restore ventilation and drainage to the normal sinus pathway diversion of obstructive tissue and polyps [4]. This minimally invasive surgical method uses endoscopic vision to create minimally manipulative surgical procedures, which can minimize trauma and facilitate faster recovery of patients, in contrast to conventional procedures works [5]. Although widely used, the effectiveness of FESS in the treatment of (CRWNP) patients is still an under-investigation topic. Varied results have been published with success rates ranging between 76%-97.5%, dependent on disease severity and surgical technique, as well as postoperative care. In addition, postsurgical remission of polyps presents a concern of concern, with rates of up to 31.33 percent recorded within an average observation of 9 Months [6,7]. The aim of the present study is to assess the clinical effectiveness of FESS in patients with (CRWNP), in terms of a symptom reduction, enhanced quality of life and postoperative recurrence. Examining these parameters, we are aiming to present the comprehensive rating of the efficiency of FESS and those factors that can have an impact on the outcomes in a long-term perspective and thus, can inform the clinical practice and the way of managing patients [8,9].

Methods

This was a cross-sectional study Conducted in Department of ENT Khyber Teaching Hospital Peshawar. From January 2023 to January 2024. Patients with the diagnosis of (CRWNP), based on the European Position Paper on Rhino sinusitis and Nasal Polyps (EPOS) criteria were included. Assessment before surgery involved a complete description of the medical history, nose endoscopy, and computed tomography (CT) par nasal sinuses. Sino-Nasal Outcome Test-22 (SNOT-22) questionnaire was used to determine the baseline

severity of symptoms and quality of life. Surgery was conducted under general anesthesia with conventional FESS procedures in an attempt to ventilate and clear the sinuses to the greatest extent possible and spare mucosa. Others required nasal saline irrigations as part of post-operative care and in some instances topical corticosteroids. Patients were reviewed after 3-, 6- and 12-months follow-up on symptom alleviation, spontaneous recurrence of the polyps, and any complications.

Inclusion Criteria

Patients aged between 18 and 65 years with medically intractable (CRWNP), proven by nasal endoscopy and CT imaging and a patient consenting to take part in the study.

Exclusion Criteria

Patients with acute rhino sinusitis, isolated sinus disease, polyps, prior Sino nasal surgery or contraindications to general anesthesia were excluded.

Approval by Ethical Committee Statement

The study was ethically accepted by the Institutional Review Board of the institution it was conducted in (approval No:IRB-no-KTH/766/11/2022). The written informed consent was signed by all participants, and it was necessary to recruit patients according to ethical standards and patient privacy in accordance with the declaration of Helsinki.

Data Collection

Sectional data collection was done through medical records, patient interview and clinical evaluations with each follow up. Demographic information, health, surgical, and postoperative data were documented carefully to provide a better analysis.

Statistical Analysis

Data analysis was done with the SPSS version 24.0 (IBM Corp., Armonk, NY, USA). Data concerning continuous variables were given as means standard deviation, and data on categorical variables were provided as frequencies and percentages. Paired t-tests were used to identify the difference between the preoperative and

Postoperative SNOT-22 scores, and p-value less than 0.05 was regarded as significant.

Results

The 75 patients mean age of 44.7212.5 years, was used. Before the surgery, SNOT-22 mean was 62.4 9.8 points, which is extreme symptomatology. Twelve months after surgery, the SNOT-22 score decreased to 28.7 +/- 7.6, which also was a significant decrease ($p < 0.001$). Nares blockage improved in 92 %, hyposmia in 63 and facial pain in 62 %. Polyp recurrence was noticed in 24 percent of the cases and the interval between the first and second recurrence was 10.2 3. reporting months. No serious adverse events (e.g., leakage of cerebrospinal fluid or loss of vision) were noted.

Table 1. Demographic Characteristics of Patients (n = 75)

Characteristic	Value
Age (years), mean \pm SD	44.7 \pm 12.5
Gender, n (%)	
Male	42 (56%)
Female	33 (44%)
Comorbidities, n (%)	
Asthma	18 (24%)
Allergic rhinitis	22 (29.3%)
Diabetes Mellitus	10 (13.3%)

Table 2. Preoperative Clinical Symptoms and SNOT-22 scores

Symptom	Preoperative n (%)	Mean \pm SD Score
Nasal obstruction	75 (100%)	8.2 \pm 1.1
Hyposmia/anosmia	60 (80%)	7.6 \pm 1.4
Facial pain/pressure	55 (73%)	6.9 \pm 1.3
Rhinorrhea	68 (90.7%)	7.1 \pm 1.2
Total SNOT-22	-	62.4 \pm 9.8

Table 3. Postoperative Symptom Improvement at 12 Months

Symptom	Improved n (%)	Mean \pm SD Score	p-value
Nasal obstruction	69 (92%)	3.2 \pm 0.9	<0.001
Hyposmia/anosmia	47 (63%)	3.5 \pm 1.1	<0.001
Facial pain/pressure	46 (62%)	3.1 \pm 1.0	<0.001
Rhinorrhea	61 (81%)	3.3 \pm 0.8	<0.001
Total SNOT-22	-	28.7 \pm 7.6	<0.001

Table 4. Recurrence and Complications

Parameter	n (%)
Polyp recurrence	18 (24%)
Mean time to recurrence (months)	10.2 \pm 3.4
Minor complications (epistaxis, crusting)	5 (6.7%)
Major complications (CSF leak, vision loss)	0 (0%)

Table 5. Effect of Surgical Extent on Outcome

Surgery Type	Patients n (%)	Mean SNOT-22 Improvement	Recurrence n (%)	p-value
Limited FESS	33 (44%)	28.2 \pm 7.5	9 (27%)	0.028
Extended FESS	42 (56%)	34.1 \pm 8.0	9 (21%)	0.028

Discussion

Functional Endoscopic Sinus Surgery (FESS) has become a basic technique in the management of chronic rhinosinusitis complicated with nasal polyp (CRWNP), which produces impressive improvements in patients with much better quality of life. Nevertheless, nasal polyps' recidivism after surgery is an issue of concern and recovery may vary between 18.2 to 40 % within 6 - 18 months after surgery [11]. This is highly variable and reflects the complexity of (CRWNP) and the necessity to develop personalized treatment strategies. The recurrence rates found in our study were similar to these findings which is 24% recurrence at 12 months post-FESS [12]. This rate can be compared with the one reported in the literature and it shows that FESS is not apparently effective in the long term, as long-lasting management approaches are necessary to ensure that the symptoms will be relieved [13]. There are a number of contributing factors to recurrence of nasal polyps after FESS. Presence of comorbid such as asthma and aspirin-exacerbated respiratory disease (AERD) has also been linked to increased recurrences [14]. We also found that asthmatics had a greater recurrence and that such comorbidities may affect surgical outcomes [15]. The scope of surgery also significantly contributes, whereby it was also found that more comprehensive surgical procedures, including extended FESS, have lower rates

of recurrence and do better in the long-term [16]. Another important issue of FESS is its effect on quality of life. In our study, there were significant improvements in SNOT-22 scores after surgery indicating improvement in nasal function, reduction in facial pain, and olfaction [17,18]. These results are in agreement with the customary findings, which showed significant positive changes in quality-of-life after FESS. It is however necessary to add that although results provide symptomatic relief, certain patients still exhibit residual symptoms and as such the continued management and education assists patients to fully grasp. Our study results in relation to surgical outcome are comparable to those of other studies. As an example, a study conducted by DeConde et al. reported 22.2% rate of recurrence of CRWNP patients who underwent a FESS. One study cited a recurrence rate of 18.2 per cent in a series of 154 patients [19,20]. These and our studies highlight the need to consider patient specific variables, including age, comorbidities and extent of surgery when planning and assessing outcomes of surgical interventions in (CRWNP) [21]. In summary, although FESS offers a valid cure to (CRWNP), nasal polyp recurrence and failure to clear symptoms in selected patients, warrants multipronged management of the disease. These are proper selection of patients, consideration of comorbidities, and the degree of surgery. P-Future study should concern itself with establishing predictive factors of recurrence and creating interventions to offset these risks so as to maintain symptom relief and quality of life enhancement over a long-term period among (CRWNP) patients [22-23].

Conclusion

fancy you are reading this article you may recognize that the olfactory sensation is the most famous one. Despite the fact that FESS leads to recurrence among a few patients, the intervention is safe and effective. With extended surgery and selecting the right patient, relapse is mitigated and victory is attained.

Limitations

Weaknesses include the study had a single center design and short follow-up of 12 months which may not represent the long-term recurrence. The sample was moderate and possible confounding factors of outcome such as the adherence to the postoperative therapy, and surgical technique variances could not be completely controlled.

Future Findings

Future studies should be attempted in multicenter format with long-term follow-up in relation to recurrence pattern over several years. Future studies are needed to understand how to integrate biologic agents, postoperative steroid regimen and individualized approaches to surgery can be developed to reduce the rate of recurrence. By pinpointing prognostic biomarkers, it is possible to further optimize patient selection and better personalize management of (CRWNP) post FESS.

Abbreviation	
FESS	Functional Endoscopic Sinus Surgery
CRWNP	Chronic Rhinosinusitis with Nasal Polyps
SNOT-22	Sino-Nasal Outcome Test-22
SD	Standard Deviation
CT	Computed Tomography
EPOS	European Position Paper on Rhinosinusitis
IRB	Institutional Review Board
CSF	Cerebrospinal Fluid
AERD	Aspirin-Exacerbated Respiratory Disease
QoL	Quality of Life
OR	Odds Ratio
RSDI	Rhinosinusitis Disability Index
E-FESS	Extended Functional Endoscopic Sinus Surgery
L-FESS	Limited Functional Endoscopic Sinus Surgery
CF	Cystic Fibrosis
GCBI	Glasgow Benefit Inventory
VAS	Visual Analog Scale
IBM SPSS	International Business Machines Statistical Package for the Social Sciences

Disclaimer: Nil **Conflict of**

Interest: Nil **Funding Disclosure:**

Nil **Authors Contribution**

Concept & Design of Study: **Imran Khan**³

Drafting: **Shakir Ullah**², **Tahreem Fatima**¹

Data Analysis: **Mushtaq Ahmad**⁴, **Muhammad Afaq**⁵

Critical Review: **Osama Nawaz**⁶, **Wasigh Ali**⁷

Final Approval : All Authors Approved the Final Version. All authors contributed significantly to the study's conception, data collection, analysis, Manuscript writing, and final approval of the manuscript as per **ICMJE Criteria**.

Ethics Statements

Studies Involving Animal Subjects

No animal studies were conducted or presented in this manuscript.

REFERENCE

1. Abdelnaseer U, El Sayed Salem A, Shawky BH, Yousef A. Assessment of Nasal Obstruction Symptoms and Pulmonary Function Following Functional Endoscopic Sinus Surgery (FESS) in Chronic Rhinosinusitis with Nasal Polyps. Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India. 2023;75:3235-41. doi: <https://doi.org/10.1007/s12070-023-03898-9>.
2. Bertlich M, Freytag S, Dombrowski T, Jurmeister P, Spiegel JL, Bertlich I, et al. Subgroups in the treatment of nasal polyposis with dupilumab: A retrospective study. Medicine. 2022;101:e31031. doi: <https://doi.org/10.1097/md.00000000000031031>.
3. Chapurin N, Khan S, Gutierrez J, Soler ZM. Economics of Medical and Surgical Management of Chronic Rhinosinusitis with Nasal Polyps: A Contemporary Review. American journal of rhinology & allergy. 2023;37:227-31. doi: <https://doi.org/10.1177/19458924221147501>.
4. Chen M, Xu M, Lei X, Zhang B. Efficacy of Endoscopic Posterior Nasal Neurectomy in Treating Eosinophilic Chronic Rhinosinusitis. ORL; journal for oto-rhino-laryngology and its related specialties. 2022;84:347-51. doi: <https://doi.org/10.1159/000519724>.
5. Deva FAL. Comparison of Conventional Nasal Douching with Corticosteroid Nasal Douching in Chronic Rhinosinusitis Patients Post Surgery. Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India. 2023;75:875-80. doi: <https://doi.org/10.1007/s12070-022-03389-3>.
6. Dharmarajan H, Falade O, Lee SE, Wang EW. Outcomes of dupilumab treatment versus endoscopic sinus surgery for chronic rhinosinusitis with nasal polyps. International forum of allergy & rhinology. 2022;12:986-95. doi: <https://doi.org/10.1002/alr.22951>.
7. Englhard AS, Ledderose GJ. Anatomical findings in patients with chronic rhinosinusitis without nasal polyps requiring revision surgery. Brazilian journal of otorhinolaryngology. 2023;89:101287. doi: <https://doi.org/10.1016/j.bjorl.2023.101287>.
8. Farrell NF, Mace JC, Sauer DA, Thomas AJ, Geltzeiler M, Detwiler KY, et al. Patient-reported sleep outcomes lack association with mucosal eosinophilia or neutrophilia in patients with chronic rhinosinusitis undergoing functional endoscopic sinus surgery. International forum of allergy & rhinology. 2021;11:784-93. doi: <https://doi.org/10.1002/alr.22678>.
9. Gupta M, Srivastava A, Kanaujia P, Kanaujia SK, Gautam HK, Saxena NS, et al. Assessment of Nasal Obstruction in Patients Undergoing Functional Endoscopic Sinus Surgery. Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India. 2023;75:3565-74. doi: <https://doi.org/10.1007/s12070-023-04019-2>.
10. Hintschich CA, Pade J, Petridis P, Hummel T. Presurgical olfactory function as an indicator of the outcome of functional endoscopic sinus surgery in chronic rhinosinusitis with nasal polyps. European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery. 2022;279:5727-33. doi: <https://doi.org/10.1007/s00405-022-07496-3>.
11. Husain S, Ramos JA, Karaf JHA, Zahedi FD, Ahmad N, Abdullah B. Efficacy of topical tranexamic acid to reduce bleeding in endoscopic sinus surgery for chronic rhinosinusitis with polyposis. European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery. 2023;280:737-41. doi: <https://doi.org/10.1007/s00405-022-07563-9>.
12. Jiang RS, Shih KH, Liang KL. Effect of Functional Endoscopic Sinus Surgery on Gustatory Function in Patients With Chronic Rhinosinusitis. Ear, nose, & throat journal. 2023;102:538-46. doi: <https://doi.org/10.1177/01455613211015754>.
13. Krishniya P, Rajpurohit P, Sharma VK, Aseri Y, Verma PC. A Prospective Correlation Study Between Computerized Tomography of Paranasal Sinuses and Nasal Endoscopy Findings in Patients of Chronic Rhinosinusitis Undergoing Functional Endoscopic Sinus Surgery. Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India. 2022;74:4706-12. doi: <https://doi.org/10.1007/s12070-021-03030-9>.
14. Loftus CA, Soler ZM, Desiato VM, Koochakzadeh S, Yoo F, Storck KA, et al. Factors impacting revision surgery in patients with chronic rhinosinusitis with nasal polyposis. International forum of allergy & rhinology. 2020;10:289-302. doi: <https://doi.org/10.1002/alr.22505>.
15. Lourijsen E, Avdeeva K, Gan KL, Pundir V, Fokkens W. Tranexamic acid for the reduction of bleeding during functional endoscopic sinus surgery. The Cochrane database of systematic reviews. 2023;2:CD012843. doi: <https://doi.org/10.1002/14651858.CD012843.pub2>.

16. Mohamed WS, El Ghonemy MT, Saber SF, Azooz KO. Risk Factors for Recurrence of Nasal Polyps after Endoscopic Sinus Surgery in Patients with Allergic Chronic Rhino Sinusitis. Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India. 2023;75:3379-85. doi: <https://doi.org/10.1007/s12070-023-03988-8>.
17. Starry A, Hardtstock F, Wilke T, Weihing J, Ultsch B, Wernitz M, et al. Epidemiology and treatment of patients with Chronic rhinosinusitis with nasal polyps in Germany-A claims data study. Allergy. 2022;77:2725-36. doi: <https://doi.org/10.1111/all.15301>.
18. Thanneru M, Lanke S, Kolavali S. The Effectiveness of Budesonide Nasal Irrigation After Endoscopic Sinus Surgery in Chronic Allergic Rhinosinusitis with Polyps. Indian journal of otolaryngology and head and neck surgery : official publication of the Association of Otolaryngologists of India. 2020;72:350-4. doi: <https://doi.org/10.1007/s12070-020-01878-x>.
19. Wang JJ, Chen CY, Liang KL, Jiang RS. Predictors of nasal bacterial culture rates in patients with chronic rhinosinusitis. European journal of clinical microbiology & infectious diseases : official publication of the European Society of Clinical Microbiology. 2020;39:711-6. doi: <https://doi.org/10.1007/s10096-019-03775-w>.
20. Wang W, Wang G, Wu T, Sun Z, Wang W, Li S. [Curative effect analysis of radical endoscopic sinus surgery for eosinophilic chronic rhinosinusitis with nasal polyps]. Lin chuang er bi yan hou tou jing wai ke za zhi = Journal of clinical otorhinolaryngology head and neck surgery. 2023;37:20-4. doi: <https://doi.org/10.13201/j.issn.2096-7993.2023.01.004>.
21. Wu YJ, Liu T, Ma JR, Wang WY, Ou SM, Zhao Y, et al. [The influence of two kinds of transnasal endoscopic surgery on the outcome status of sinus cavity in patients with eosinophilic chronic rhinosinusitis with nasal polyps]. Zhonghua er bi yan hou tou jing wai ke za zhi = Chinese journal of otorhinolaryngology head and neck surgery. 2022;57:1450-6. doi: <https://doi.org/10.3760/cma.j.cn115330-20220905-00540-1>.
22. Zhang L, Zhang Y, Gao Y, Wang K, Lou H, Meng Y, et al. Long-term outcomes of different endoscopic sinus surgery in recurrent chronic rhinosinusitis with nasal polyps and asthma. Rhinology. 2020;58:126-35. doi: <https://doi.org/10.4193/Rhin19.184>.
23. Zhang M, Ryan PJ, Shashinder S. Efficacy of Absorbable Steroid-Impregnated Nasal Packing in Functional Endoscopic Sinus Surgery for Chronic Rhinosinusitis: A Systematic Review. The Laryngoscope. 2021;131:1704-14. doi: <https://doi.org/10.1002/lary.29350>.



Licensing and Copyright Statement

All articles published in the **Pakistan Journal of Advances in Medicine and Medical Study (PJAMMR)** are licensed under the terms of the **Creative Commons Attribution-NonCommercial-4.0, International License (CC BY-NC 4.0)**. This license permits Non-Commercial Use, distribution, and reproduction in any medium, provided the original author and source are properly cited. Commercial use of the content is not permitted, without prior permission from the **Author(s) 2025** the journal.

[This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.](#)